

Talk at MTNS 2014, Groningen, The Netherlands

by Salma Kuhlmann, Universität Konstanz, Germany

Application of the Archimedean
Positivstellensatz to locally multiplicatively
convex real algebras

July 10, 2014

THE K -MOMENT PROBLEM

Let $A := \mathbb{R}[\underline{X}] := \mathbb{R}[X_1, \dots, X_n]$ be the algebra of polynomials in n variables with real coefficients and $L : A \rightarrow \mathbb{R}$ a real valued linear functional.

The K -moment problem

Given $\emptyset \neq K \subseteq \mathbb{R}^n$, when is L representable as an integral with respect to a positive Borel measure, i.e.

$$L(f) = \int_K f d\mu, \quad \forall f \in \mathbb{R}[\underline{X}],$$

where μ is supported on K ?

THE K -MOMENT PROBLEM

Haviland, 1936

Such a measure exists if and only if $L(\text{Psd}(K)) \subseteq [0, \infty)$, where $\text{Psd}(K) := \{f \in A : f(x) \geq 0 \quad \forall x \in K\}$.

THE K -MOMENT PROBLEM

Haviland, 1936

Such a measure exists if and only if $L(\text{Psd}(K)) \subseteq [0, \infty)$, where $\text{Psd}(K) := \{f \in A : f(x) \geq 0 \quad \forall x \in K\}$.

Scheiderer, 1999

Except for a few cases, checking $L(\text{Psd}(K)) \subseteq [0, \infty)$ is not a finite procedure, i.e. $\text{Psd}(K)$ usually is not *finitely generated*.

DEFINITIONS

- ▶ $M \subseteq A$ is a **quadratic module**:

- ▶ M is a cone:

$$0, 1 \in M, \quad M + M \subseteq M \text{ and } [0, \infty) \cdot M \subseteq M.$$

- ▶ $\forall f \in A \quad f^2 M \subseteq M.$

DEFINITIONS

- ▶ $M \subseteq A$ is a **quadratic module**:

- ▶ M is a cone:

$$0, 1 \in M, \quad M + M \subseteq M \text{ and } [0, \infty) \cdot M \subseteq M.$$

- ▶ $\forall f \in A \quad f^2 M \subseteq M.$

- ▶ M is **Archimedean**:

$$\forall f \in A \quad \exists N \geq 1 \quad N + f \in M.$$

DEFINITIONS

- ▶ $M \subseteq A$ is a **quadratic module**:

- ▶ M is a cone:

$$0, 1 \in M, \quad M + M \subseteq M \text{ and } [0, \infty) \cdot M \subseteq M.$$

- ▶ $\forall f \in A \quad f^2 M \subseteq M.$

- ▶ M is **Archimedean**:

$$\forall f \in A \quad \exists N \geq 1 \quad N + f \in M.$$

- ▶ A quadratic module T with $T \cdot T \subseteq T$ is called a **preordering**.

DEFINITIONS

- ▶ $M \subseteq A$ is a **quadratic module**:

- ▶ M is a cone:

$$0, 1 \in M, \quad M + M \subseteq M \text{ and } [0, \infty) \cdot M \subseteq M.$$

- ▶ $\forall f \in A \quad f^2 M \subseteq M.$

- ▶ M is **Archimedean**:

$$\forall f \in A \quad \exists N \geq 1 \quad N + f \in M.$$

- ▶ A quadratic module T with $T \cdot T \subseteq T$ is called a **preordering**.
- ▶ Let $S \subset A$; M_S (resp. T_S):= The smallest quadratic module (resp. preordering) containing S .

DEFINITIONS

- ▶ $M \subseteq A$ is a **quadratic module**:

- ▶ M is a cone:

$$0, 1 \in M, \quad M + M \subseteq M \text{ and } [0, \infty) \cdot M \subseteq M.$$

- ▶ $\forall f \in A \quad f^2 M \subseteq M.$

- ▶ M is **Archimedean**:

$$\forall f \in A \quad \exists N \geq 1 \quad N + f \in M.$$

- ▶ A quadratic module T with $T \cdot T \subseteq T$ is called a **preordering**.
- ▶ Let $S \subset A$; M_S (resp. T_S):= The smallest quadratic module (resp. preordering) containing S .
- ▶ M (or T) is **finitely generated**, if $M = M_S$ (or $T = T_S$) for some finite S .

DEFINITIONS

- ▶ $M \subseteq A$ is a **quadratic module**:
 - ▶ M is a cone:

$$0, 1 \in M, \quad M + M \subseteq M \text{ and } [0, \infty) \cdot M \subseteq M.$$

- ▶ $\forall f \in A \quad f^2 M \subseteq M.$
- ▶ M is **Archimedean**:

$$\forall f \in A \quad \exists N \geq 1 \quad N + f \in M.$$

- ▶ A quadratic module T with $T \cdot T \subseteq T$ is called a **preordering**.
- ▶ Let $S \subset A$; M_S (resp. T_S):= The smallest quadratic module (resp. preordering) containing S .
- ▶ M (or T) is **finitely generated**, if $M = M_S$ (or $T = T_S$) for some finite S .
- ▶ $S \subset A$:

$$\mathcal{K}_S := \{x \in \mathbb{R}^n : f(x) \geq 0 \quad \forall f \in S\}.$$

CLASSICAL SOLUTIONS

Schmüdgen, 1991

If S is finite and \mathcal{K}_S is compact, then

$$L(T_S) \subseteq [0, \infty) \Rightarrow L(\text{Psd}(\mathcal{K}_S)) \subseteq [0, \infty).$$

CLASSICAL SOLUTIONS

Schmüdgen, 1991

If S is finite and \mathcal{K}_S is compact, then

$$L(T_S) \subseteq [0, \infty) \Rightarrow L(\text{Psd}(\mathcal{K}_S)) \subseteq [0, \infty).$$

Putinar, 1993

If S is finite and M_S is Archimedean, then

$$L(M_S) \subseteq [0, \infty) \Rightarrow L(\text{Psd}(\mathcal{K}_S)) \subseteq [0, \infty).$$

CLASSICAL SOLUTIONS

Schmüdgen, 1991

If S is finite and \mathcal{K}_S is compact, then

$$L(T_S) \subseteq [0, \infty) \Rightarrow L(\text{Psd}(\mathcal{K}_S)) \subseteq [0, \infty).$$

Putinar, 1993

If S is finite and M_S is Archimedean, then

$$L(M_S) \subseteq [0, \infty) \Rightarrow L(\text{Psd}(\mathcal{K}_S)) \subseteq [0, \infty).$$

Since T_S and M_S are finitely generated, Haviland's Theorem is effectively applicable to them.

TOPOLOGICAL INTERPRETATION

φ := The *finest locally convex* topology on A .

TOPOLOGICAL INTERPRETATION

φ := The *finest locally convex* topology on A .

Schmüdgen:

$$\text{Psd}(\mathcal{K}_S) = \overline{T_S}^{\varphi}$$

TOPOLOGICAL INTERPRETATION

$\varphi :=$ The *finest locally convex* topology on A .

Schmüdgen:

$$\text{Psd}(\mathcal{K}_S) = \overline{T}_S^\varphi$$

Putinar:

$$\text{Psd}(\mathcal{K}_S) = \overline{M}_S^\varphi$$

TOPOLOGICAL INTERPRETATION

$\varphi :=$ The *finest locally convex* topology on A .

Schmüdgen:

$$\text{Psd}(\mathcal{K}_S) = \overline{T}_S^\varphi$$

Putinar:

$$\Rightarrow \text{Psd}(K) = \overline{C}^T$$

$$\text{Psd}(\mathcal{K}_S) = \overline{M}_S^\varphi$$

TOPOLOGICAL INTERPRETATION

$\varphi :=$ The *finest locally convex* topology on A .

Schmüdgen:

$$\text{Psd}(\mathcal{K}_S) = \overline{T}_S^\varphi$$

Putinar:

$$\Rightarrow \text{Psd}(K) = \overline{C}^\tau$$

$$\text{Psd}(\mathcal{K}_S) = \overline{M}_S^\varphi$$

- ▶ For a locally convex topology τ on A ,
- ▶ C is a convex cone of A ,
- ▶ and K is a closed subset of \mathbb{R}^n .

EXAMPLE

1. Replace φ by $\|\cdot\|_K$ -topology, where $K = [-1, 1]^n$ and

$$\|f\|_K := \sup_{x \in K} |f(x)|.$$

$$\text{Stone-Weierstrass} \Rightarrow \text{Psd}(K) = \overline{\sum \mathbb{R}[\underline{X}]^2}^{\|\cdot\|_K}.$$

EXAMPLE

1. Replace φ by $\|\cdot\|_K$ -topology, where $K = [-1, 1]^n$ and

$$\|f\|_K := \sup_{x \in K} |f(x)|.$$

Stone-Weierstrass $\Rightarrow \text{Psd}(K) = \overline{\sum \mathbb{R}[\underline{X}]^2}^{\|\cdot\|_K}$.

2. Replace φ by $\|\cdot\|_1$ -topology, $K = [-1, 1]^n$ where

$$\left\| \sum_{\alpha} f_{\alpha} \underline{X}^{\alpha} \right\|_1 := \sum_{\alpha} |f_{\alpha}|.$$

Berg *et al.* $\Rightarrow \text{Psd}(K) = \overline{\sum \mathbb{R}[\underline{X}]^2}^{\|\cdot\|_1}$.

EXAMPLE

In term of moments:

If L is a $\|\cdot\|_K$ or $\|\cdot\|_1$ - continuous positive semidefinite functional, then there exists a Borel measure μ on $[-1, 1]^n$ such that

$$\forall f \in \mathbb{R}[\underline{X}] \quad L(f) = \int_{[-1,1]^n} f \, d\mu.$$

GENERAL SETTINGS:

Now, let A be a unital commutative \mathbb{R} -algebra and $\mathcal{X}(A) := \text{Hom}_{\mathbb{R}}(A, \mathbb{R}) \subseteq \mathbb{R}^A$ endowed with the product topology.

GENERAL SETTINGS:

Now, let A be a unital commutative \mathbb{R} -algebra and $\mathcal{X}(A) := \text{Hom}_{\mathbb{R}}(A, \mathbb{R}) \subseteq \mathbb{R}^A$ endowed with the product topology. To every $a \in A$ we associate a map

$$\begin{aligned} \hat{a} : \mathcal{X}(A) &\rightarrow \mathbb{R} \\ \alpha &\mapsto \alpha(a) \end{aligned}$$

GENERAL SETTINGS:

Now, let A be a unital commutative \mathbb{R} -algebra and $\mathcal{X}(A) := \text{Hom}_{\mathbb{R}}(A, \mathbb{R}) \subseteq \mathbb{R}^A$ endowed with the product topology. To every $a \in A$ we associate a map

$$\begin{aligned} \hat{a} : \mathcal{X}(A) &\rightarrow \mathbb{R} \\ \alpha &\mapsto \alpha(a) \end{aligned}$$

$$K \subseteq \mathcal{X}(A) \longrightarrow \text{Psd}(K) := \{a \in A : \hat{a} \geq 0 \text{ on } K\},$$

GENERAL SETTINGS:

Now, let A be a unital commutative \mathbb{R} -algebra and $\mathcal{X}(A) := \text{Hom}_{\mathbb{R}}(A, \mathbb{R}) \subseteq \mathbb{R}^A$ endowed with the product topology. To every $a \in A$ we associate a map

$$\begin{aligned} \hat{a} : \mathcal{X}(A) &\rightarrow \mathbb{R} \\ \alpha &\mapsto \alpha(a) \end{aligned}$$

$$\begin{aligned} K \subseteq \mathcal{X}(A) &\longrightarrow \text{Psd}(K) := \{a \in A : \hat{a} \geq 0 \text{ on } K\}, \\ S \subseteq A &\longrightarrow \mathcal{K}_S := \{\alpha \in \mathcal{X}(A) : \alpha(S) \subseteq [0, \infty)\}. \end{aligned}$$

GENERAL SETTINGS:

Now, let A be a unital commutative \mathbb{R} -algebra and $\mathcal{X}(A) := \text{Hom}_{\mathbb{R}}(A, \mathbb{R}) \subseteq \mathbb{R}^A$ endowed with the product topology. To every $a \in A$ we associate a map

$$\begin{aligned} \hat{a} : \mathcal{X}(A) &\rightarrow \mathbb{R} \\ \alpha &\mapsto \alpha(a) \end{aligned}$$

$$\begin{aligned} K \subseteq \mathcal{X}(A) &\longrightarrow \text{Psd}(K) := \{a \in A : \hat{a} \geq 0 \text{ on } K\}, \\ S \subseteq A &\longrightarrow \mathcal{K}_S := \{\alpha \in \mathcal{X}(A) : \alpha(S) \subseteq [0, \infty)\}. \end{aligned}$$

A $\sum A^{2d}$ -module ($d \geq 1$ an integer), is a cone $C \subseteq A$ such that

$$\forall a \in A \quad a^{2d} \cdot C \subseteq C.$$

GENERAL SETTINGS:

Now, let A be a unital commutative \mathbb{R} -algebra and $\mathcal{X}(A) := \text{Hom}_{\mathbb{R}}(A, \mathbb{R}) \subseteq \mathbb{R}^A$ endowed with the product topology. To every $a \in A$ we associate a map

$$\begin{aligned} \hat{a} : \mathcal{X}(A) &\rightarrow \mathbb{R} \\ \alpha &\mapsto \alpha(a) \end{aligned}$$

$$\begin{aligned} K \subseteq \mathcal{X}(A) &\longrightarrow \text{Psd}(K) := \{a \in A : \hat{a} \geq 0 \text{ on } K\}, \\ S \subseteq A &\longrightarrow \mathcal{K}_S := \{\alpha \in \mathcal{X}(A) : \alpha(S) \subseteq [0, \infty)\}. \end{aligned}$$

A $\sum A^{2d}$ -module ($d \geq 1$ an integer), is a cone $C \subseteq A$ such that

$$\forall a \in A \quad a^{2d} \cdot C \subseteq C.$$

T. Jacobi's Theorem, 2001

Let C be an Archimedean $\sum A^{2d}$ -module of A . Then for each $a \in A$

$$\hat{a} > 0 \text{ on } \mathcal{K}_C \Rightarrow a \in C.$$

SEMINORMED ALGEBRAS

A map $\rho : A \longrightarrow [0, \infty)$ is called a **seminorm** if

$$1 \quad \forall a \in A \quad \forall r \in \mathbb{R} \quad \rho(ra) = |r|\rho(a),$$

$$2 \quad \forall a, b \in A \quad \rho(a + b) \leq \rho(a) + \rho(b);$$

ρ is **submultiplicative** if

$$3 \quad \forall a, b \in A \quad \rho(ab) \leq \rho(a)\rho(b).$$

The pair (A, ρ) is called a seminormed algebra.

SEMINORMED ALGEBRAS

A map $\rho : A \longrightarrow [0, \infty)$ is called a **seminorm** if

$$1 \quad \forall a \in A \quad \forall r \in \mathbb{R} \quad \rho(ra) = |r|\rho(a),$$

$$2 \quad \forall a, b \in A \quad \rho(a + b) \leq \rho(a) + \rho(b);$$

ρ is **submultiplicative** if

$$3 \quad \forall a, b \in A \quad \rho(ab) \leq \rho(a)\rho(b).$$

The pair (A, ρ) is called a seminormed algebra.

The **Gelfand spectrum** of (A, ρ) :

$$\text{sp}(\rho) := \{ \alpha \in \mathcal{X}(A) : \alpha \text{ is } \rho\text{-continuous} \}$$

SEMINORMED ALGEBRAS

A map $\rho : A \rightarrow [0, \infty)$ is called a **seminorm** if

$$1 \quad \forall a \in A \quad \forall r \in \mathbb{R} \quad \rho(ra) = |r|\rho(a),$$

$$2 \quad \forall a, b \in A \quad \rho(a + b) \leq \rho(a) + \rho(b);$$

ρ is **submultiplicative** if

$$3 \quad \forall a, b \in A \quad \rho(ab) \leq \rho(a)\rho(b).$$

The pair (A, ρ) is called a seminormed algebra.

The **Gelfand spectrum** of (A, ρ) :

$$\begin{aligned} \text{sp}(\rho) &:= \{ \alpha \in \mathcal{X}(A) : \alpha \text{ is } \rho\text{-continuous} \} \\ &= \{ \alpha \in \mathcal{X}(A) : |\alpha(a)| \leq \rho(a) \quad \forall a \in A \} \end{aligned}$$

SEMINORMED ALGEBRAS

A map $\rho : A \longrightarrow [0, \infty)$ is called a **seminorm** if

$$1 \quad \forall a \in A \quad \forall r \in \mathbb{R} \quad \rho(ra) = |r|\rho(a),$$

$$2 \quad \forall a, b \in A \quad \rho(a + b) \leq \rho(a) + \rho(b);$$

ρ is **submultiplicative** if

$$3 \quad \forall a, b \in A \quad \rho(ab) \leq \rho(a)\rho(b).$$

The pair (A, ρ) is called a seminormed algebra.

The **Gelfand spectrum** of (A, ρ) :

$$\begin{aligned} \text{sp}(\rho) &:= \{ \alpha \in \mathcal{X}(A) : \alpha \text{ is } \rho\text{-continuous} \} \\ &= \{ \alpha \in \mathcal{X}(A) : |\alpha(a)| \leq \rho(a) \quad \forall a \in A \} \\ &\subseteq \prod_{a \in A} [-\rho(a), \rho(a)], \end{aligned}$$

is a compact Hausdorff space.

SEMINORMED ALGEBRAS

$I_\rho := \{a \in A : \rho(a) = 0\}$ is an *ideal* of A and

$$\begin{aligned}\bar{\rho} : \bar{A} = A/I_\rho &\rightarrow [0, \infty) \\ \bar{a} &\mapsto \rho(a)\end{aligned}$$

induces a norm on \bar{A} which admits a *completion* $(\tilde{A}, \tilde{\rho})$ and

$$\mathfrak{sp}(\rho) \sim \mathfrak{sp}(\tilde{\rho}).$$

SEMINORMED ALGEBRAS

$I_\rho := \{a \in A : \rho(a) = 0\}$ is an *ideal* of A and

$$\begin{aligned}\bar{\rho} : \bar{A} = A/I_\rho &\rightarrow [0, \infty) \\ \bar{a} &\mapsto \rho(a)\end{aligned}$$

induces a norm on \bar{A} which admits a *completion* $(\tilde{A}, \tilde{\rho})$ and

$$\mathfrak{sp}(\rho) \sim \mathfrak{sp}(\tilde{\rho}).$$

Lemma

Let $(B, \|\cdot\|)$ be a Banach algebra, $a \in B$, $r > \|a\|$ and $k \geq 1$ an integer. Then there exist $b \in B$ such that $b^k = r + a$. Thus any $\sum B^{2d}$ -module is archimedean and any $\alpha \in \mathcal{X}(B)$ is continuous.

SEMINORMED ALGEBRAS

$I_\rho := \{a \in A : \rho(a) = 0\}$ is an *ideal* of A and

$$\begin{aligned} \bar{\rho} : \bar{A} = A/I_\rho &\rightarrow [0, \infty) \\ \bar{a} &\mapsto \rho(a) \end{aligned}$$

induces a norm on \bar{A} which admits a *completion* $(\tilde{A}, \tilde{\rho})$ and

$$\mathfrak{sp}(\rho) \sim \mathfrak{sp}(\tilde{\rho}).$$

Lemma

Let $(B, \|\cdot\|)$ be a Banach algebra, $a \in B$, $r > \|a\|$ and $k \geq 1$ an integer. Then there exist $b \in B$ such that $b^k = r + a$. Thus any $\sum B^{2d}$ -module is archimedean and any $\alpha \in \mathcal{X}(B)$ is continuous.

Proof.

The Taylor expansion of $(1+t)^{\frac{1}{k}} = \sum_{i=0}^{\infty} \lambda_i t^i$ converges absolutely for $|t| < 1$. Now set $t := \frac{a}{r}$. □

MAIN RESULT

Theorem 1

Let (A, ρ) be a seminormed \mathbb{R} -algebra, $d \geq 1$ an integer, $C \subseteq A$ a $\sum A^{2d}$ -module. Then

$$\overline{C}^\rho = \text{Psd}(\mathcal{K}_C \cap \text{sp}(\rho)).$$

MAIN RESULT

Proof.

$C \subseteq \text{Psd}(\mathcal{K}_C \cap \mathfrak{sp}(\rho)) = \bigcap_{\alpha \in \mathcal{K}_C \cap \mathfrak{sp}(\rho)} \alpha^{-1}([0, \infty))$ which is closed.

Therefore $\overline{C}^\rho \subseteq \text{Psd}(\mathcal{K}_C \cap \mathfrak{sp}(\rho))$.

MAIN RESULT

Proof.

$C \subseteq \text{Psd}(\mathcal{K}_C \cap \mathfrak{sp}(\rho)) = \bigcap_{\alpha \in \mathcal{K}_C \cap \mathfrak{sp}(\rho)} \alpha^{-1}([0, \infty))$ which is closed.

Therefore $\bar{C}^\rho \subseteq \text{Psd}(\mathcal{K}_C \cap \mathfrak{sp}(\rho))$. For the other inclusion, set $\tilde{C} :=$ The closure of the image of C in $(\tilde{A}, \tilde{\rho})$.

MAIN RESULT

Proof.

$C \subseteq \text{Psd}(\mathcal{K}_C \cap \mathfrak{sp}(\rho)) = \bigcap_{\alpha \in \mathcal{K}_C \cap \mathfrak{sp}(\rho)} \alpha^{-1}([0, \infty))$ which is closed.

Therefore $\bar{C}^\rho \subseteq \text{Psd}(\mathcal{K}_C \cap \mathfrak{sp}(\rho))$. For the other inclusion, set $\tilde{C} :=$ The closure of the image of C in $(\tilde{A}, \tilde{\rho})$. \tilde{C} is a $\sum \tilde{A}^{2d}$ -module of \tilde{A} . So \tilde{C} is **Archimedean**.

MAIN RESULT

Proof.

$C \subseteq \text{Psd}(\mathcal{K}_C \cap \mathfrak{sp}(\rho)) = \bigcap_{\alpha \in \mathcal{K}_C \cap \mathfrak{sp}(\rho)} \alpha^{-1}([0, \infty))$ which is closed.

Therefore $\overline{C}^\rho \subseteq \text{Psd}(\mathcal{K}_C \cap \mathfrak{sp}(\rho))$. For the other inclusion, set $\tilde{C} :=$ The closure of the image of C in $(\tilde{A}, \tilde{\rho})$. \tilde{C} is a $\sum \tilde{A}^{2d}$ -module of \tilde{A} . So \tilde{C} is **Archimedean**.

Take $b \in \text{Psd}(\mathcal{K}_C \cap \mathfrak{sp}(\rho))$ with image \tilde{b} in \tilde{A} . For any $\alpha \in \mathcal{K}_{\tilde{C}}$ we have $0 \leq \alpha(\tilde{b}) = \alpha|_A(b)$, so $\forall n \geq 1 \forall \alpha \in \mathcal{K}_{\tilde{C}} \quad \alpha(\frac{1}{n} + \tilde{b}) > 0$.

MAIN RESULT

Proof.

$C \subseteq \text{Psd}(\mathcal{K}_C \cap \mathfrak{sp}(\rho)) = \bigcap_{\alpha \in \mathcal{K}_C \cap \mathfrak{sp}(\rho)} \alpha^{-1}([0, \infty))$ which is closed.

Therefore $\overline{C}^{\rho} \subseteq \text{Psd}(\mathcal{K}_C \cap \mathfrak{sp}(\rho))$. For the other inclusion, set $\tilde{C} :=$ The closure of the image of C in $(\tilde{A}, \tilde{\rho})$. \tilde{C} is a $\sum \tilde{A}^{2d}$ -module of \tilde{A} . So \tilde{C} is **Archimedean**.

Take $b \in \text{Psd}(\mathcal{K}_C \cap \mathfrak{sp}(\rho))$ with image \tilde{b} in \tilde{A} . For any $\alpha \in \mathcal{K}_{\tilde{C}}$ we have $0 \leq \alpha(\tilde{b}) = \alpha|_A(b)$, so $\forall n \geq 1 \forall \alpha \in \mathcal{K}_{\tilde{C}} \quad \alpha(\frac{1}{n} + \tilde{b}) > 0$. By **T. Jacobi's Theorem**, $\tilde{b} + \frac{1}{n} \in \tilde{C}$. Letting $n \rightarrow \infty$, $\tilde{b} \in \tilde{C}$, and hence $b \in \overline{C}^{\rho}$. \square

CORRESPONDING MOMENT PROBLEM

Corollary

Let $L : A \longrightarrow \mathbb{R}$ be a ρ -continuous linear functional. If $L(C) \subseteq [0, \infty)$ then there exists a unique Radon measure μ on $\mathcal{K}_C \cap \mathfrak{sp}(\rho)$ such that

$$L(a) = \int \hat{a} d\mu, \quad \forall a \in A.$$

LOCALLY MULTIPLICATIVELY CONVEX TOPOLOGIES

Let \mathcal{F} be a family of submultiplicative seminorms on A . The family \mathcal{F} induces a locally convex topology $\tau_{\mathcal{F}}$ on A such that $(A, \tau_{\mathcal{F}})$ is a topological algebra.

LOCALLY MULTIPLICATIVELY CONVEX TOPOLOGIES

Let \mathcal{F} be a family of submultiplicative seminorms on A . The family \mathcal{F} induces a locally convex topology $\tau_{\mathcal{F}}$ on A such that $(A, \tau_{\mathcal{F}})$ is a topological algebra.

A topology τ is said to be *locally multiplicatively convex (lmc)* if $\tau = \tau_{\mathcal{F}}$ for some family \mathcal{F} of submultiplicative seminorms on A .

LOCALLY MULTIPLICATIVELY CONVEX TOPOLOGIES

Let \mathcal{F} be a family of submultiplicative seminorms on A . The family \mathcal{F} induces a locally convex topology $\tau_{\mathcal{F}}$ on A such that $(A, \tau_{\mathcal{F}})$ is a topological algebra.

A topology τ is said to be *locally multiplicatively convex (lmc)* if $\tau = \tau_{\mathcal{F}}$ for some family \mathcal{F} of submultiplicative seminorms on A .

Proposition

If \mathcal{F} is saturated then $\mathfrak{sp}(\tau_{\mathcal{F}}) = \bigcup_{\rho \in \mathcal{F}} \mathfrak{sp}(\rho)$.

CLOSURES AND MOMENTS IN LMC TOPOLOGIES

Theorem 2

Let τ be an lmc topology on A , $d \geq 1$ an integer, C a $\sum A^{2d}$ -module. Then

$$\overline{C}^\tau = \text{Psd}(\mathcal{K}_C \cap \text{sp}(\tau)).$$

CLOSURES AND MOMENTS IN LMC TOPOLOGIES

Theorem 2

Let τ be an lmc topology on A , $d \geq 1$ an integer, C a $\sum A^{2d}$ -module. Then

$$\overline{C}^\tau = \text{Psd}(\mathcal{K}_C \cap \text{sp}(\tau)).$$

Corollary

Let $L : A \rightarrow \mathbb{R}$ be a τ -continuous functional with $L(C) \subseteq [0, \infty)$. Then there exists a unique Radon measure μ on $\mathcal{K}_C \cap \text{sp}(\tau)$ such that

$$L(a) = \int \hat{a} d\mu, \quad \forall a \in A.$$

SCHMÜDGEN'S RESULT

Schmüdgen, 1978

Let η be the finest lmc topology on A and $d \geq 1$. Then

$$\overline{\sum A^{2d}}^{\eta} = \text{Psd}(\mathcal{X}(A)).$$

INVOLUTIVE \mathbb{C} -ALGEBRAS

Let $(A, \rho, *)$ be a seminormed \mathbb{C} -algebra equipped with an involution $*$.

INVOLUTIVE \mathbb{C} -ALGEBRAS

Let $(A, \rho, *)$ be a seminormed \mathbb{C} -algebra equipped with an involution $*$.

- ▶ $\mathcal{X}_*(A) := \{\alpha : A \longrightarrow \mathbb{C} : \alpha \text{ is a } *\text{-algebra homomorphism}\},$
- ▶ $\mathfrak{sp}_*(\rho) := \{\alpha \in \mathcal{X}_*(A) : \alpha \text{ is } \rho\text{-continuous}\},$
- ▶ $H(A) := \{a \in A : a^* = a\}.$

INVOLUTIVE \mathbb{C} -ALGEBRAS

Let $(A, \rho, *)$ be a seminormed \mathbb{C} -algebra equipped with an involution $*$.

- ▶ $\mathcal{X}_*(A) := \{\alpha : A \longrightarrow \mathbb{C} : \alpha \text{ is a } *\text{-algebra homomorphism}\},$
- ▶ $\mathfrak{sp}_*(\rho) := \{\alpha \in \mathcal{X}_*(A) : \alpha \text{ is } \rho\text{-continuous}\},$
- ▶ $H(A) := \{a \in A : a^* = a\}.$

Corollary

Let $C \subseteq H(A)$ be a $\sum H(A)^{2d}$ -module of $H(A)$. Let $L : A \longrightarrow \mathbb{C}$ be a ρ -continuous $*$ -functional such that $L(C) \subseteq [0, \infty)$. Then there exists a unique Radon measure μ on $\mathcal{K}_C \cap \mathfrak{sp}_*(\rho)$ such that

$$L(a) = \int \hat{a} d\mu, \quad \forall a \in A.$$

BERG-MASERICK

Let $(S, 1, *)$ be a commutative unitary $*$ -semigroup.

An *absolute value* on S is a map $\phi : S \rightarrow [0, \infty)$ such that

1. $\phi(1) \geq 1$,
2. $\forall s, t \in S, \phi(st) \leq \phi(s)\phi(t)$,
3. $\forall s \in S \quad \phi(s^*) = \phi(s)$.

BERG-MASERICK

Let $(S, 1, *)$ be a commutative unitary $*$ -semigroup.

An *absolute value* on S is a map $\phi : S \rightarrow [0, \infty)$ such that

1. $\phi(1) \geq 1$,
2. $\forall s, t \in S, \phi(st) \leq \phi(s)\phi(t)$,
3. $\forall s \in S \quad \phi(s^*) = \phi(s)$.

The map $\|\cdot\|_\phi$ on $\mathbb{C}[S]$ defined by $\|\sum_s f_s s\|_\phi = \sum_s |f_s| \phi(s)$ is a submultiplicative seminorm on $\mathbb{C}[S]$.

BERG-MASERICK

Let $(S, 1, *)$ be a commutative unitary $*$ -semigroup.

An *absolute value* on S is a map $\phi : S \rightarrow [0, \infty)$ such that

1. $\phi(1) \geq 1$,
2. $\forall s, t \in S, \phi(st) \leq \phi(s)\phi(t)$,
3. $\forall s \in S \quad \phi(s^*) = \phi(s)$.

The map $\|\cdot\|_\phi$ on $\mathbb{C}[S]$ defined by $\|\sum_s f_s s\|_\phi = \sum_s |f_s| \phi(s)$ is a submultiplicative seminorm on $\mathbb{C}[S]$.

Berg-Maserick, 1984

If $L : \mathbb{C}[S] \rightarrow \mathbb{C}$ is an $*$ -functional such that

$L(\sum H(\mathbb{C}[S])^{2d}) \subseteq [0, \infty)$ and $\exists c > 0 \forall s \in S \quad |L(s)| \leq c\phi(s)$. Then there exists a unique Radon measure μ on $\mathfrak{sp}_*(\|\cdot\|_\phi)$ such that

$$L(f) = \int \hat{f} d\mu \quad \forall f \in \mathbb{C}[S].$$

Thank you